

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 December 2000 (28.12.2000)

PCT

(10) International Publication Number
WO 00/78320 A1

- (51) International Patent Classification⁷: A61K 31/70, 35/78, 38/00, 59/00, A23L 1/30, 2/00 (74) Agents: LAPPIN, Mark, G. et al.; McDermott, Well & Emery, 28 State Street, Boston, MA 02109 (US).
- (21) International Application Number: PCT/US00/40267 (81) Designated States (*national*): AU, CA, JP, MX.
- (22) International Filing Date: 21 June 2000 (21.06.2000) (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 09/338,021 22 June 1999 (22.06.1999) US
- (71) Applicant: JOINT JUICE INCORPORATED [US/US]; 3727 Buchanan Street, San Francisco, CA 94123 (US).
- (72) Inventor: STONE, Kevin, R.; 1 Throckmorton Lane, Mill Valley, CA 94941 (US).

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 00/78320 A1

(54) Title: CARTILAGE ENHANCING FOOD SUPPLEMENTS AND METHODS OF PREPARING THE SAME

(57) Abstract: A food supplement, either in the form of a snack bar or a beverage, which contains one or more cartilage enhancing supplements is provided. The cartilage supplements include chondroitin, glucosamine, and hyaluronic acid. The food supplement may additionally be fortified with cetyl myristoleate. The beverage is a mixture of a juice drink base which may include a water-based fruit flavored juice prepared using a pasteurization process at a relatively high temperature and a cartilage supplement solution which includes a cartilage supplement prepared at a relatively low temperature. The beverage may be carbonated, non-carbonated or concentrated. The preferred cartilage supplement is glucosamine, preferably associated with a counter ion, more preferably as glucosamine HCl.

CARTILAGE ENHANCING FOOD SUPPLEMENTS AND METHODS OF PREPARING THE SAME

5 FIELD OF THE INVENTION

The present invention relates to food supplements, such for example, snack bars and beverages which are fortified with one or more cartilage supplements.

BACKGROUND

10 Nutritional bars and energy drinks are convenient nutritional supplements, particularly for those persons too busy to eat regular meals and for hikers, cyclists, runners or other athletes who need prepackaged, ready-to-eat, high-energy snacks while they are exercising. Such bars and drinks are also convenient nutritional supplements for the elderly who need prepackaged, ready-to-eat snacks. Additionally, such food
15 supplements can supply consumers with the necessary vitamins and minerals specified in the recommended daily allowances provided by the U.S. government.

By way of example, U.S. Patent No. 4,543,262 discloses a high protein, low or no lactose, vitamin and mineral fortified, nutritionally-balanced snack bar. Additionally, U.S. Patent No. 3,814,819 teaches a protein-fortified food bar composed of several baked
20 crisp wafers layered on top of the other with a creamy filling between them. The creamy filling contains added vitamins, providing twenty-five percent (25%) of the recommended daily allowance of vitamins and minerals. U.S. Patent No. 4,152,462 teaches a highly nutritious protein and vitamin enriched food bar, having a marshmallow base. Lastly, U.S. Patent No. 3,901,799 discloses a high protein chocolate bar. Caseinate and peanut
25 butter are added to a mixture of chocolate and cocoa butter. Vitamins compatible with the ingredients, it is disclosed, can be added to the snack bar.

Drinks formulations and methods for preparing them have also been developed. Energy enhancement drinks and processes have been developed such as in U.S. Patent No. 3,894,148, which relates to nutritional and exercise therapy to maximize the storage
30 of glycogen in muscle tissue. Protein fortification drinks have been described in U.S. Patent No. 4,309,417. Finally, carbohydrate and electrolyte drinks have been described in U.S. Patent No. 4,312,856 and 4,322,407.

It has been long recognized that dietary cartilage supplements are effective in reducing the symptoms of joint pain. (Drovani, Clinical Therapeutics, (1980)). An amino-acid complex combining glutamine with glucosamine sulfate is the constituent component used by the body to make cartilage and connective tissue, which cushion and lubricate the joints in the body. To date, more than 6,000 patients have been studied in 20 clinical trials. These studies have all reached the conclusion that glucosamine sulfate supplements relieve pain and stimulate healing in osteoarthritis patients. In fact, the World Health Organization has officially classified glucosamine sulfate as a slow-acting drug for the treatment of osteoarthritis. Typically, glucosamine sulfate is taken in the form of a pill or a powder.

Chondroitin sulfate is another compound widely sold as an agent for the treatment of the symptoms of joint pain. Its healing properties as a dietary supplement, however, while still effective, have proven in two studies to be lower than the healing properties of glucosamine sulfate. Chondroitin sulfate is also taken in the form of a pill or a powder.

Hyaluronic acid is a polysaccharide which forms a major component of the gel-like substance found in the connective tissue of mammals. Structurally, it is comprised of a repeating disaccharide consisting of N-acetyl-D-glucosamine and D-glucuronic acid. Functionally, it serves as a lubricant and shock absorbent in mammalian joints. Hyaluronic acid is also taken in the form of a pill or a powder and is thought to be effective for the treatment of joint pain.

Cetyl myristoleate (CM) is a newly recognized agent that is potentially useful for the treatment of joint pain. CM is an ester of a fatty acid, the building blocks of fats and oils. CM is produced by combining the fatty acid myristolic acid with cetyl alcohol, a long-chain alcohol. CM appears to function in three ways in the body. First, it shows an anti-inflammatory effect. Second, it appears to act as a lubricant for joints. Third, CM functions as an immune system modulator.

Doctors have reported that significant improvement in patients with osteoarthritis who were taking CM. In 1996, in a one-month multi-center clinical study involving 431 patients with various forms of arthritis, sixty-three percent (63%) of those taking CM showed improvement compared to fourteen percent (14%) in the control group. CM is typically taken orally in the form of an oil.

Konjac flour is a soluble dietary fiber that is similar in structure and function to pectin and typically is used as a thickener, emulsifier, gelling agent, film former and stabilizer. Glucomannan, the main constituent of Konjac flour, is a slightly branched hydrocolloidal polysaccharide having B1-4 linked subunits of glucose and mannose and
5 having a molecular weight ranging between 200,000 and 2,000,000 daltons. Acetyl groups, located every 9-19 subunits along the glucomannan backbone, help solubilize the molecule. In addition to being a food agent, glucomannan has been tested on humans, principally as a means for lowering serum cholesterol, bile acid level and serum triglyceride. Studies indicate that glucomannan may affect glucose tolerance and glucose
10 absorption.

Stevia (*Stevia rebaudiana bectoni*) is a natural, non-caloric sweet-tasting plant that is typically used in medical applications for inhibiting fat absorption and for lowering blood pressure as well as in the food industry as a non-caloric sweetening agent. Stevioside is the component of Stevia that gives the plant its sweetness. As a sugar
15 substitute, it is available as a concentrated liquid, crushed leaf, or concentrated white powder. Often, individuals who do not tolerate sugar or other sweeteners can use Stevia. Medicinally, studies indicate that Stevia helps regulate the pancreas and may help stabilize blood sugar levels within the body. Stevia is also indicated as a cardi tonic, for obesity, to reduce stomach acidity, to reduce gas, for hypotension and to help lower uric
20 acid levels. Research has also indicated that Stevia may help reduce bacteria.

While carbohydrates, proteins and fats are all important in the human diet, carbohydrates are particularly important for athletic performance. Carbohydrates are a well-known source of energy which are readily absorbed by the body. For example, marathon runners and other athletes typically "carbo-load" the day before a race by eating
25 large amounts of carbohydrates. Moreover, athletes in endurance events need a source of energy which is readily absorbable by the body in order to replace the diminishing stores of glucose and glycogen that occur during the event. Lastly, athletes typically consume large quantities of carbohydrates immediately following a race in order to replenish glycogen levels depleted by the event. Thus, the energy source provided by
30 carbohydrates is important to athletes before, during, and after the race.

Typically, carbohydrates range between complex carbohydrates and simple sugars. Structurally, these carbohydrates differ in the number of sugars in the molecule and in the degree of branching. Functionally, they differ by how readily the body can absorb them and process them to derive energy. Thus, the correct ratio of the different types of carbohydrates can supply short-term, mid-term, and long-term supplies of energy to the body.

During athletic events, particularly endurance events such as marathon running, triathalons and long distance cycling, athletes can deplete much, if not all, of their glycogen stores. It is therefore important that athletes replenish their depleted stores of glycogen from a source of carbohydrates. Typically, the cellular machinery used to convert glucose to glycogen is most efficient in the several hours immediately following the athletic event, the so-called recovery period. In addition to depleting their glycogen stores, athletes can cause temporary, and sometimes permanent, damage to the joints of their bodies. Typically, they experience this damage as pain and stiffness in their joints. In some forms, the present invention provides a quality source of carbohydrates, which is important during the recovery period. The invention also provide supplements for the carbohydrate source. In accordance with the invention, the supplements include one or more cartilage supplements which also aid in recovery by reducing joint pain and stiffness, and/or Konjac flour and/or Stevia supplements. The supplements are provided alone or in combination with CM.

It is therefore one object of this invention to provide a formula which provides a nutritional snack together with cartilage supplements which address cartilage dysfunction, and/or a Konjac flour supplement and/or a Stevia supplement.

It is another object of the invention to provide a formula which provides a nutritional snack together with cartilage supplements which address cartilage dysfunction, and/or a Konjac flour supplement and/or a Stevia supplement, in combination with CM.

It is yet another object of this invention to provide a nutritional snack which aids athletes in recovering from athletic events.

Another object of the invention to provide a formula for a beverage including one or more cartilage supplements which address cartilage dysfunction.

It is a further object of the invention to provide a beverage that is source of quality carbohydrates and one or more cartilage supplements.

Yet another object of the invention is to provide a method of preparing a beverage containing one or more cartilage supplements.

5

SUMMARY OF THE INVENTION

The present invention provides a food supplement, for human and/or animal consumption and methods for making such food supplements. The food supplement can be either in the form of a beverage or a snack bar, or preparation for making such
10 beverages or snack bars. The food supplement of the present invention is fortified with one or more cartilage supplements including glucosamine sulfate, chondroitin sulfate and hyaluronic acid. The food supplements of the invention may also be fortified with Konjac flour and/or Stevia supplements either alone or in combination with the cartilage supplements. The food supplement, as disclosed by the invention, can be optionally
15 further fortified with cetyl myristoleate (CM).

In one preferred embodiment, the present invention provides a high protein, nutritional snack bar that is fortified with one or more cartilage, Konjac flour and/or Stevia supplements, and which can also be fortified with CM. The snack bar has pleasing textural and taste characteristics. In another preferred embodiment, the present invention
20 provides a beverage which is fortified with one or more cartilage, Konjac flour and/or Stevia supplements, and which can be additionally fortified with CM.

All of the compounds used to fortify the food supplement, according to the present invention, are safe for human consumption. Moreover, in addition to the cartilage supplement, any salt, counter ion or derivative of the supplement which is safe for human
25 consumption, and preferably dissociates to the cartilage supplement, either in water or after ingestion, is suitable for use in the present invention. Suitable cartilage supplements are hyaluronic acid, chondroitin, and preferably glucosamine. For chondroitin, sulfate is the preferred counter ion. Where the cartilage supplement is glucosamine, suitable derivatives include but are not limited to, glucosamine n-alkyl sulfonate, and n-acetyl
30 glucosamine. Preferable counter ions for glucosamine include, but are not limited to, phosphoric, sulfate, and, most preferably, hydrochloride. Glucosamine hydrochloride

provides improved taste to the nutritional supplements, and thereby minimizes the need for flavorings used to mask the strong flavors characteristic of other cartilage supplements.

In one preferred embodiment, the food supplement containing the cartilage, Konjac
5 flour and/or Stevia supplements, alone or in combination with CM, is a nutritional snack bar. The process of preparing the snack bar, according to the present invention, comprises melting a confectioner's material, which is a solid at ambient temperature. The melted confectioner's material is admixed with the major liquid ingredients and with the major and minor dry ingredients. If the snack bar, according to the present invention, is
10 to be fortified with CM, the CM is first admixed with the major liquid ingredients. Admixing the minor dry compounds with the major liquid ingredients prior to admixing with the confectioner's material minimizes localized concentrations of dry ingredients.

The above mixture is extruded in an extruder typical of extruders known to those skilled in the art. The extruded material or extrudate is then cut to a desired size. The
15 snack bar can be coated on the surface with a melted confectioner's coating material and/or a topping to enhance taste and visual appeal.

In another preferred embodiment, the above-referenced snack bar is modified to include one or more reservoirs, each reservoir containing cartilage, Konjac flour and/or Stevia supplements or mixtures of two or more of such supplements. Within each
20 reservoir, the supplements can be further combined with CM. By way of example, the snack bar can contain a substance such as chocolate or carob, which is molded with one or more internal void regions or reservoirs.

In another preferred embodiment, the cartilage, supplements and/or Konjac flour and/or Stevia supplements and/or the CM are provided in the form of a beverage.
25 Beverages suitable for use in the present invention include naturally or artificially flavored fruit juices, commercially available sports drinks such as GATORADE™ and POWERADE™ beverages, and commercially available nutritionally-balanced beverages such as ENSURE™ beverage.

In a preferred method of making beverages of the invention, a "cold mix" process
30 is used. Pursuant to that process, in a preferred form as noted above, the juice drink base

is most preferably a fruit flavored juice. However, the juice drink base could alternatively be unflavored water.

A beverage for human or animal consumption is made from a mixture of a juice drink base and a cartilage supplement-containing solution. The juice drink base is a water-based solution, and is preferably a flavored (and most preferably a naturally or artificially fruit-flavored) juice. In one embodiment, the juice drink base is prepared using a pasteurization process at a relatively high temperature. The high temperature is naturally or artificially sufficient to kill bacteria. The cartilage supplement-containing solution is a water-based solution containing one or more cartilage supplements (such as glucosamine, chondroitin, hyaluronic acid, or mixtures thereof) prepared at a relatively low temperature. Preferably, the low temperature is in a range that minimizes inactivation of the cartilage supplement. The beverage may be carbonated or non-carbonated; furthermore, the beverage can be in a concentrated form for later dilution by the consumer or ready to drink.

In yet another embodiment of the invention there is provided a method of treating cartilage dysfunction in a human by administering a cartilage enhancing beverage formulated according to the above aspects of invention.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Commercially available confectioner's coating material is used in the present invention. The compositions of typical confectioner's coating materials which can be used in the present invention, are disclosed in cookie and cracker technology, *AVI Publishing Co., Westport, Conn., page 176, table 45 (1968)*. These materials are generally solid at room temperature, but melt and can be poured when heated. The confectioner's material in the present invention preferably has melting points below 140° F and more preferably below 105° F.

Confectioner's material with higher melting or pouring points can present several difficulties. For example, the higher the melting point of the coating material, the greater is the risk that the heated coating material will begin to set or partially solidify when admixed with the other ingredients that are preferably maintained at room temperature. Premature hardening of the coating material can cause a lack of uniform distribution of

the confectioner's material within the snack bar. As a result, any ingredients added to the confectioner's material may also not be uniformly distributed throughout the product. This is particularly true of flavoring ingredients which may be added to the melted confectioner's material.

5 In the present invention, the confectioner's coating material may be cocoa butter-based or alternatively may be a compound coating, which is made from hardened vegetable oils. Examples of hardened vegetable oils used for compound coatings include cottonseed, coconut, soybean, palm and peanut oils. These hardened vegetable oils are mixed with sugar and are the principle ingredients in the coating material. Compound
10 coatings may be peanut flavored, fruit flavored, chocolate flavored, vanilla flavored, coconut flavored or flavored with other commercially available flavorings.

 The snack bar according to the present invention, can be fortified with protein from several sources of edible protein, either alone or in combination. Exemplary sources include whey powder, carob powder, soy lecithin, peanut flour, wheat proteins such as
15 wheat germ and caseinates such as calcium, potassium and sodium caseinates.

 Exemplary sources of carbohydrates suitable for the snack bar of the present invention include malted cereal syrup from corn, barley and brown rice; maltodextran; fructose; high fructose corn syrup; date paste; sucrose; brown sugar and mixtures thereof. Exemplary sources of complex carbohydrates include those derived from cereal grains
20 such as oats, rice, barley, and corn. Preferably, the snack bar according to the present invention, contains at least one sugar and at least one cereal grain as the sources of carbohydrates.

 In the present invention, corn syrup, preferably high fructose corn syrup, is used as a source of carbohydrates. The corn syrup gives the snack a moist chewy texture,
25 provides a source of sweetness to the snack bar, and aids in distributing the dry ingredients. Additionally, the corn syrup together with the confectioner's coating material, binds together the other ingredients such as the protein source ingredients and the cereal grain ingredients.

 Fats have the highest source of energy per unit weight, approximately twice that of
30 proteins or carbohydrates. In addition to the confectioner's coating material, other sources of fat suitable for use in the present invention include flavoring such as chocolate,

cocoa, and coconut and at least one partially hydrogenated vegetable oil such as soybean, cottonseed, corn and palm oil.

The above-listed ingredients are categorized based on the main nutritional component that each contributes to the snack bar. It is, however, useful to note that many
5 of the ingredients may be sources of two or more nutritional components. For example, whey powder, peanut flour, and wheat germ each contain substantial amounts of carbohydrates, proteins, and fats. By varying the ingredients in the snack bar, one can maintain the caloric distribution disclosed by the present invention.

Several additional ingredients are preferably added to the snack bar according to
10 the present invention. For example, the snack bar may be topped with conventional toppings such as crushed nuts, granola and the like. Additionally, although shortenings other than the confectioner's coating are not necessary, it is preferable to include a wetting agent to facilitate mixing and binding of the dry ingredients with the confectioner's coating and corn syrup and to enhance the moisture and chew capability of
15 the snack. Suitable wetting agents for use in the present invention include molasses, honey, and the partially hydrogenated vegetable oils.

The nutritional snacks according to the present invention, are made by first melting the confectioner's coating material by heating it to its melting point or to within about ten degrees (10°) Fahrenheit above its melting point. Next, the liquid ingredients which
20 comprise the corn syrup, the melted confectioner's coating material, and the optional wetting agent are mixed together. If the snack bar is to be fortified with cetyl myristoleate (CM), the CM is added to the liquid ingredients. The liquid ingredients are mixed to substantial homogeneity.

Once the liquid ingredients are thoroughly mixed, the minor dry ingredient are
25 mixed into the liquid ingredients. The minor dry ingredients include the chondroitin sulfate, the glucosamine sulfate, the hyaluronic acid and the optional salt. The minor dry ingredients are mixed with the liquids in order to avoid localized high concentrations of the minor dry ingredients.

Once the minor dry ingredients are mixed to substantial homogeneity with the
30 liquid ingredients, the major dry ingredients are then admixed to the liquid ingredients and the minor dry ingredients. This mixture is mixed to substantial homogeneity. Major dry

ingredients include the high content carbohydrates such as sugars and cereal grains and the high content proteins such as the whey protein, soy protein, peanut protein and caseinate.

Flavoring ingredients such as cocoa or coconut can be added to the mixture.

- 5 When adding flavoring ingredients, it is preferable to add them to the liquid ingredients to avoid localized concentrations within the snack bar. Once all the ingredients are mixed to substantial homogeneity, the mixture is transferred to a conventional confectionery-type bar extruder. The mixture is forced through a die of the extruder to form the extrudate. The die may be any shape known to those skilled in the art, but is preferably rectangular.
- 10 The extrusion is done at room temperature.

After extrusion, the extrudate is cut to the desired size. The cut extrudate may then be coated by, for example, dipping with a melted confectioner's coating material. The bar is then chilled and may then be topped with a topping such as granola or ground nuts. The bar is then packaged for shipping or storage.

- 15 In another preferred embodiment, the cartilage supplements, either alone or in combination with the CM, are supplied in the form of a beverage. By way of non-limiting example, beverages suitable for use in the present invention include water or fruit juices such as apple juice, orange juice, grapefruit juice, cranberry juice, pineapple juice, grape juice, and mixtures thereof. Other beverages suitable for use in the present invention
- 20 include commercially available sports drinks such as GATORADE™ and POWERADE™ beverages. Generally, these sports drinks provide calories substantially only in the form of carbohydrates and provide electrolyte replacements, which are thought to aid in recovery after exercise. Lastly, commercially available nutritionally balanced beverages, such for example ENSURE™ beverage, are suitable for use in the present invention.
- 25 These nutritionally balanced beverages generally provide carbohydrates, protein, fat, vitamins and minerals to consumers and often serve as meal replacements.

- In accordance with the invention, all three cartilage supplements, either alone or in combination, are included in beverages, such as the above-identified beverages. Where CM is included, it is important to note, however, that CM is an oily-like substance and,
- 30 therefore, tends not to be readily miscible with the aqueous based beverages. Accordingly, when CM is added to a beverage according to the present invention, it is

first emulsified. Emulsifying agents and methods known to those skilled in the art are suitable for use in the present invention. The emulsifying agent chosen is fit for human consumption. Suitable emulsifying agents include gum arabic and gelatin. The emulsions are prepared by shaking together the two liquids or by adding one phase drop-wise to the other with some form of agitation, such as irradiation by high intensity ultrasonic waves.

In a preferred method, the cartilage enhanced beverages made according to the present invention are made using a multi-step process. As a first step, a juice drink base is prepared. Water, flavorings, and preservatives are mixed together in predetermined amounts. The solution is heat pasteurized, and cooled to ambient temperatures. The juice base may be prepared and stored for later use, or used immediately in the production of the final beverage products.

As a second step, a second solution of "cartilage supplement solution", containing the cartilage supplement, and, if and as desired, additional flavorings and carbohydrates, is prepared at temperatures below those used in a heat pasteurization process, preferably but not necessarily, at ambient temperatures or lower. As a third step, the drink base is combined with the cartilage supplement solution at ambient temperatures. The combined solution is adjusted for acidity, and preferably, but not necessarily, chilled to temperatures below ambient and above freezing, and carbonated. The multi-step method insures that the cartilage supplement is not affected by the preparation and/or bottling process. As an alternative, the second and third steps may be combined, where the ingredients for the cartilage supplement solution are added directly to the juice drink base, and pH adjusted, and if desired, carbonated. That combined second and third step is performed, preferably but not necessarily, at temperatures below those used in a heat pasteurization process.

In another embodiment, a beverage concentrate is prepared. Juice base, sweetener, water, preservatives and, optionally, vitamins, and/or minerals, are mixed in a mixing tank and pasteurized. The resulting product is transferred from the mixing tank to the packaging through transfer lines. The product cools in the transfer lines as it flows from the mixing tank to the packaging. The product temperature is lowered by radiational cooling of the product flowing through the transfer lines, or optionally, the lines may be chilled.

Preferably, the cartilage supplement solution is introduced into the liquid stream at a point in transfer lines, preferably at a distance from the mixing tank, where the temperature of the flowing liquid is lower than the temperature in the mixing tank. This allows the cartilage solution to mix into the product at temperatures below pasteurization temperature and as the beverage solution is cooling continuously in transit. Once in the packaging, the containers are sealed, and sprayed or doused with cool water to lower the temperature of the final product to approximately ambient temperature.

In this embodiment, the total residence time of the cartilage solution at elevated temperatures is minimal, thereby minimizing any heat inactivation of the cartilage supplement. For example, the product temperature may be close to 195°F at the mix tank. As the fluid progresses through the transfer lines, the point where the glucosamine is added may be 150°F or less. As the solution passes to the packaging it is continually cooled so that at the point of packaging, the solution temperature at the packaging point may be approximately 100°F to 80°F. After packaging, the final product is cooled to ambient temperature. Thus, in this example, the total residence time of glucosamine near pasteurization temperatures may be only about 30 seconds while undergoing continuous cooling.

The heat pasteurization process for the juice base is dependent on both the temperature employed and the residence time of the solution at that temperature. Generally, and within a range, the lower the temperature used, the longer the solution should be processed to insure adequate microbial control. Pasteurization parameters can range from a low of about 165°F for 3 minutes to about 200°F for less than 40 seconds. The preferable parameters are $195 \pm 4^\circ\text{F}$ for 42 ± 4 seconds.

In an alternate embodiment, the cartilage supplement may be added directly to the juice base and other ingredients at or near pasteurization temperatures. In yet another embodiment, the cartilage supplement may be added to the mix tank with the juice base and other ingredients, and the product may be pasteurized using pasteurization processes that do not include heat processing, such as, but not limited to, filter sterilization or radiation sterilization.

The juice bases, pre-carbonated product, and concentrated beverage solutions are measured for Brix (a measure of soluble solids), and titratable acidity (TA) and adjusted

as necessary, using water and citric acid, respectively. The pre-carbonated ready to drink product and concentrated beverage product are also pH adjusted just before carbonation and packaging, respectively.

The beverages produced by these various embodiment include, but are not limited to ready to drink carbonated beverages, ready to drink non-carbonated beverages, and concentrated beverage mixes, i.e., mixes that are diluted to drinking concentration (with water, juice, carbonated water, and the like) by the consumer.

EXAMPLE 1

10 In this example, a snack bar with a surface coating is prepared. The ingredients and relative amounts are shown in Table 1.

The snack bar of the present invention is made by melting the confectioner's peanut butter material and mixing in a conventional mixer, the liquid components which comprise the high fructose corn syrup and the molasses.

15 The second step is to add the minor dry ingredients comprising the salt, chondroitin sulfate, glucosamine sulfate and the hyaluronic acid, to the liquid mixture. The mixture is mixed to substantial homogeneity.

The third step in the process is adding the major dry ingredients to the mixed ingredients of the second process step. The major dry ingredients include the sugars, whey protein, rice flour and soy protein. The mixture is then mixed to substantial homogeneity. The mixture is then fed into a conventional extruding machine.

20 The mixture is extruded at room temperature. As the mixture is extruded, the extrudate is cut into individual serving sizes of about sixty (60) grams. The cut pieces are then coated with approximately ten (10) grams of melted chocolate confectioner's material. The bar is then allowed to cool or is chilled and is wrapped for shipping and storage. The final bar is approximately seventy (70) grams and has about three hundred (300) calories.

EXAMPLE 2

30 A nutritional bar with a core, a surface coating and a topping is prepared. The bar is further fortified with CM. The ingredients and relative amounts are shown in Table 2.

The snack bar is made by melting chocolate confectioner's coating material. The corn syrup, vegetable oil and CM are next added to the melted confectioner's coating.

Next, the minor dry ingredients, as set forth in Example 1, are mixed with the liquids, along with the flavorings. The mixture is mixed to substantial homogeneity.

5 To this mixture, the major dry ingredients are then added. These include the bran, oats, barley, fructose, caseinate, and the wheat germ. The mixture is then mixed to substantial homogeneity.

The mixture is then processed as set forth in Example 1. The bar is then coated with approximately ten (10) grams of melted chocolate confectioner's coating and then
10 topped with crushed nuts. The bar is then packaged for shipping and storage.

EXAMPLE 3

The beverage of the present invention containing cartilage supplements is produced as follows:

15 First, a volume of GATORADE is aliquoted into a conventional fluid mixer. Hyaluronic acid and/or, glucosamine sulfate, and/or chondroitin sulfate are next added to the solution to a final concentration of four percent (4%) (weight to volume) of each component. The solution is next mixed until the added ingredients become solubilized. The solution is then aseptically filled into individual serving size bottles of approximately
20 five hundred milliliters (500 ml).

EXAMPLE 4

The beverage of the present invention containing a cartilage supplement and CM is produced as follows:

25 First, CM is emulsified with any known emulsifying agent suitable for human consumption. Next, a volume of ENSURE is aliquoted into a conventional fluid mixer. The emulsified CM is added to a final concentration of two percent (2%) (volume to volume). Next, the hyaluronic acid, glucosamine sulfate, and chondroitin sulfate are added to the solution to a final concentration of four percent (4%) (weight to volume).
30 The solution is mixed until the added cartilage supplements become solubilized. The

solution is then aseptically filled into individual serving size bottles of approximately five hundred milliliters (500 ml).

EXAMPLES 5-7

5 *Preparation of the Juice Drink Base.*

Juice drink base solutions containing the ingredients and proportions shown in Table 3 were prepared and stored for later use formulating the finished beverage products.

For each flavor, the above ingredients were mixed together using 90% of the water. The Brix, titratable acidity and pH were adjusted, using the remaining 10% of the water. The resultant solution was pasteurized by heating the solution to 195°F and maintaining it at that temperature for approximately 40 seconds. Then the solution was cooled to ambient temperature to complete preparation of the drink base. In these examples of the juice drink base, the Brix, and titratable acidity values were adjusted to the ranges shown in Table 4. These values may vary by about ± 5 percent. The juice base was then stored for later use in formulating the final products.

Examples 8-10

Preparation of Ready to Drink Beverages: Ready to drink cartilage enhancing supplement-enhancing beverages were prepared at ambient temperature using the ingredients (and proportions) shown in Table 5.

Initially, sodium benzoate was added to 50% of the total water, and mixed at ambient temperature until the sodium benzoate was dissolved. That solution was maintained under continuous agitation. The glucosamine HCl, citric acid, Juice Drink Base and high fructose corn syrup were each added separately to the solution and mixed thoroughly. Ninety percent (90%) of the remaining water was then added and the solution was mixed for 15 minutes. The brix, titratable acidity and pH were adjusted using the remaining water, and if necessary, using additional citric acid. The resulting solution was then carbonated in a proportioner (1 + 4) at 40°F to 3.0 volumes. The resulting beverage was then ready for bottling.

30 The acceptable ranges for Brix, and TA in the beverages before and after carbonation, and pH after carbonation are shown in Table 6:

Examples 11-13.

Preparation of Sugar Free Ready to Drink Beverages: Sugar free (diet) ready to drink cartilage enhancing supplement-enhancing beverages were prepared at ambient temperature using the ingredients (and proportions) shown in Table 7.

5 Initially, sodium benzoate was added to 50% of the total water, and mixed at ambient temperature until the sodium benzoate was dissolved. That solution was maintained under continuous agitation. The glucosamine HCl, citric acid, aspartame, acesulfame K, and Juice Base were each added separately to the solution and mixed thoroughly. Ninety percent (90%) of the remaining water was then added and the solution
10 was mixed for 15 minutes. The brix, titratable acidity and pH were adjusted using the remaining water, and if necessary, using additional citric acid. The resulting solution was then carbonated in a proportioner (1+4), to 3.0 volumes, and further adjusting the pH. The resulting beverage was then ready for bottling.

Brix, TA, and pH of the beverages before and after carbonation are measured.

15 Table 8 shows the acceptable ranges after carbonation.

Examples 14-17.

Preparation of Concentrated Juice Bases. Concentrated cartilage enhancing supplement beverages were prepared using the ingredients (and proportions) shown in
20 Table 9.

Preparation of Concentrated Beverage: Initially, 50% of the water was pumped into the mixing tank. Sodium benzoate was added to the water and thoroughly mixed. Citric acid, sweetener (high fructose corn syrup or artificial sweeteners) and Juice Base were added to the solution and thoroughly mixed. Brix and TA were adjusted using a measured
25 amount of the remaining water and citric acid, respectively. The concentrated juice bases were heat pasteurized at approximately 195°F for about 42 seconds.

The solution was pumped through transfer lines to the packaging area. Glucosamine HCl (dissolved in a measured amount of water) was fed into the fluid stream in the transfer lines. Brix, TA and pH were tested on the finished product, and the batch parameters were
30 adjusted appropriately, as shown in Table 10. The packages were sealed, and rinsed with

water at ambient temperature. Yield: approximately 7,060 two ounce packages, depending on flavor.

Those of skill in the art will recognize that the invention may be embodied in other specific forms without departing from the spirit of essential characteristics thereof. The presently described embodiments are therefore to be considered in all respects as illustrative and not restrictive, the scope of the invention being indicated by the appended claims rather than by the foregoing description. All changes that come within the meaning and range of the equivalency of the claims are therefore intended to be embraced therein.

10

Table 1. Snack Bar Formulation

INGREDIENTS	TOTAL WEIGHT (%)
Whey Protein	24.5
Sugars	8.0
Rice Flour	16.0
Soy Protein	6.0
Chondroitin Sulfate	1.5
Glucosamine Sulfate	1.5
Hyaluronic Acid	1.5
Salt	0.5
Corn Syrup	28.5
Molasses	4.0
Confectioner's Peanut Butter	8.0
TOTAL	100.0

Table 2. Nutritional Bar with Topping

INGREDIENTS	WEIGHT PERCENTAGE
Bran	2.5
Oat	10.0
Barley	4.0
Fructose	6.0
Caseinate	11.0
Flavorings	4.0
Wheat Germ	13.0
Chondroitin Sulfate	1.5
Glucosamine Sulfate	1.5
Hyaluronic Acid	1.5
Salt	0.5
Corn Syrup	25.0
Partially Hydrogenated Vegetable Oil	1.0
Cetyl Myristoleate (CM)	1.5
Chocolate Confectioner's Compound Coating	18.5
TOTAL	100.0

Table 3. Juice Base Formulations

INGREDIENTS	Example 5 Orange		Example 6 Lemon		Example 7 Tropical	
	Wt %	lb.	Wt. %	lb.	Wt %	lb.
Water	20.6419	47.30	2.93	6.72	16.7984	41.69
Ascorbic Acid	0.8505	1.95	1.75	4.00	0.8500	2.11
Natural Juicy Citrus Accord	1.2430	2.85	2.55	5.84	2.0922	5.19
Sodium Benzoate	0.1000	0.23	0.10	0.23	0.1000	0.25
Potassium Sorbate	0.1000	0.23	0.10	0.23	0.1000	0.25
Orange Juice Concentrate	65.0274	149.01	--	--	55.8683	138.64
Natural Tangerine Flavor	7.3270	16.79	--	--	2.6153	6.49
Natural Orange Flavor Powder	2.6168	6.00	--	--	2.0922	5.19
Natural Orange Flavor Emulsion	2.0934	4.80	--	--	--	--
Lemon Juice Concentrate	--	--	69.95	160.18	--	--
Natural Lemon Flavor, WONF	--	--	19.40	44.43	--	--
Natural Lemon Extract Flavor	--	--	3.22	7.38	--	--
Mango Concentrate	--	--	--	--	12.1609	30.18
Natural Mango Flavor	--	--	--	--	0.9480	2.35
Passionfruit Concentrate	--	--	--	--	5.2305	12.98
Natural Red Color, liquid	--	--	--	--	1.1442	2.84
TOTAL (gallons)	100.00	229.16 (25)	100.00	229.01 (24)	100.00	248.16 (25)

Table 4. Brix, and TA* values for Juice Base

	Orange	Lemon	Tropical
Brix	42.6	33.3	42.7
TA	3.13	29.72	3.63

*TA = titratable acidity; acceptable ranges are within $\pm 5\%$.

Table 5. Ready to Drink Beverages

	Example 8		Example 9		Example 10	
INGREDIENTS	Orange		Lemon		Tropical	
	Wt %	lb.	Wt. %	lb.	Wt %	lb.
Water	39.81	591.40	62.45	872.16	39.63	589.43
High Fructose Corn Syrup	42.02	624.24	28.39	396.43	41.96	624.32
Glucosamine HCl	2.38	35.36	2.41	33.69	2.38	35.37
Citric Acid	2.21	32.90	0.00	0.00	2.46	36.60
Sodium Benzoate	0.23	3.37	0.24	3.31	0.23	3.37
Tangerine Orange Base (Example 5)	13.35	198.38	--	--	--	--
Lemon Juice Base (Example 6)	--	--	6.51	91.22	--	--
Tropical Juice Base (Example 7)	--	--			13.34	198.52
TOTAL (gallons)	100.00	1485.65 (150)	100.00	1396.81 (150)	100.00	1487.61 (150)

Table 6. Brix, TA* and pH Acceptable Ranges Before and After Carbonation

	Orange		Lemon		Tropical	
	Before Carb.	After Carb.	Before Carb.	After Carb.	Before Carb.	After Carb.
Brix	43.1° ±5%	9.2-10.4°	27.0° ±5%	7.0-8.6°	43.3° ±5%	9.2-10.4
TA	2.62 ±5%	0.53-0.58	1.95 ±5%	0.55-0.57	3.63 ±5%	0.55-0.66
pH	--	3.48-3.56	--	3.49-3.56	--	3.34-3.55

*TA = titratable acidity

Table 7: Sugar Free Ready to Drink Beverages

	Example 11		Example 12		Example 13	
INGREDIENTS	Orange Diet		Lemon Diet		Tropical Diet	
	Wt %	lb.	Wt. %	lb.	Wt %	lb.
Water	81.795	1215.64	89.981	1256.91	81.560	1213.74
Aspartame	0.025	0.37	0.267	0.37	0.025	0.37
Glucosamine HCl	2.380	35.36	2.407	33.63	2.377	35.37
Citric Acid	2.214	32.90	0.805	11.25	2.460	36.60
Sodium Benzoate	0.227	3.37	0.237	3.31	0.226	3.37
Acesulfame K	0.012	0.18	0.013	0.18	0.012	0.18
Tangerine Orange Base (Example 5)	13.347	198.38	--	--	--	--
Lemon Juice Base (Example 6)	--	--	6.530	91.22	--	--
Tropical Juice Base (Example 7)	--	--	--	--	13.340	198.52
TOTAL (gallons)	100.00	1486.20 (150)	100.00	1396.87 (150)	100.00	1488.15 (150)

Table 8. Target Values and Ranges for pH, Brix and TA* for Sugar Free Beverages

	Diet Orange	Diet Lemon	Diet Tropical
	After Carbonation	After Carbonation	After Carbonation
Brix Range	2-4°	2-4°	2-4°
TA Range	0.68-0.73	0.65-0.70	0.82-0.87
pH Range	2.75-2.95	2.70-2.90	2.80-2.95

*TA=titratable acidity

Table 9: Concentrated Juice Beverages

	Example 14		Example 15		Example 16		Example 17	
INGREDIENTS	Orange		Lemon		Diet Lemon		Tropical	
	Wt %	lb.	Wt. %	lb.	Wt.%	lb.	Wt %	lb.
Water	17.401	88.24	47.978	511.34	86.430	921.22	17.106	85.59
High Fructose Corn Syrup	57.736	24.58	38.504	410.37	--	--	57.735	26.40
Glucosamine HCl	3.158	34.16	3.157	33.65	3.157	33.65	3.158	34.26
Citric Acid	3.052	33.01	1.093	11.65	1.094	11.66	3.387	36.74
Sodium Benzoate	0.317	3.43	0.326	3.48	0.326	3.48	0.317	3.44
Aspartame	--	--	--	--	0.035	0.38	--	--
Acesulfame K	--	--	--	--	0.018	0.19	--	--
Tangerine Orange Base (Example 5)	18.336	98.38	--	--	--	--	--	--
Lemon Juice Base (Example 6)	--	--	8.942	95.29	8.940	95.29	--	--
Tropical Juice Base (Example 7)	--	--	--	--	--	--	18.297	98.52
TOTAL	100.00	081.80	100.00	1065.78	100.00	1065.87	100.00	084.94

Table 10. Acceptable Ranges for pH, Brix and TA* for Concentrate Beverages

	Orange	Diet Lemon	Lemon	Tropical
Brix	55-59°	7-11°	35-39°	55-59°
TA	4.10-4.17	3.85-3.93	3.75-3.83	4.55-4.60
pH	2.25-2.40	2.25-2.35	2.30-2.45	2.20-2.35

*TA = titratable acidity

What is claimed is:

- 1 1. A beverage for human consumption comprising:
2 a mixture of a juice drink base and a cartilage supplement-containing solution
3 wherein:
4 A. said juice drink base is a water-based solution comprising - juice, and
5 prepared using a pasteurization process at a relatively high temperature, and
6 B. said cartilage-supplement containing solution is a water-based
7 solution comprising one or more cartilage supplements and prepared at a relatively
8 low temperature.
- 1 2. A beverage according to claim 1 wherein said mixture has a pH in the range of 2.2-
2 4.6.
- 1 3. A beverage according to claim 1 wherein said cartilage supplement is glucosamine.
- 1 4. A beverage according to claim 1 wherein said cartilage supplement is chondroitin.
- 1 5. A beverage according to claim 3 wherein said relatively high temperature is above
2 190°F, and said relatively low temperature is below 150°F.
- 1 6. A beverage according to claim 1 wherein said cartilage supplement is a salt or
2 derivative of the cartilage supplement.
- 1 7. A beverage according to claim 6 wherein the cartilage supplement is glucosamine
2 hydrochloride.

- 1 8. A beverage according to claim 6 wherein in said relatively high temperature is
2 above 190°F, and said relatively low temperature is below 150°F.
- 1 9. A beverage according to claim 1 wherein juice is fruit flavored.
- 1 10. A beverage according to claim 9 wherein said fruit flavored juice comprises a fruit
2 juice.
- 1 11. A beverage according to claim 10 wherein said fruit-flavored juice is a fruit juice
2 concentrate.
- 1 12. A beverage according to claim 1 wherein said mixture is carbonated.
- 1 13. A beverage according to claim 1 wherein said mixture is not carbonated
- 1 14. A beverage according to claim 1 wherein said mixture is effected at a relatively
2 low mix temperature.
- 1 15. A beverage according to claim 14 wherein said low mix temperature is
2 approximately 40-160°F.
- 1 16. A beverage according to claim 3 wherein said cartilage supplements are selected
2 from the group consisting of glucosamine salts, glucosamine derivatives, chondroitin,
3 chondroitin sulfate, hyaluronic acid, and combinations thereof.
- 1 17. A beverage according claim 15 wherein the cartilage supplement is glucosamine
2 hydrochloride.

- 1 18. A method of preparing a beverage for human consumption, comprising the steps of
2 A. providing a juice drink base, said juice drink base being a water-
3 based solution comprising a fruit-flavored juice prepared using a pasteurization
4 process at a relatively high temperature;
5 B. providing a water-based cartilage supplement solution comprising a
6 cartilage supplement, said cartilage supplement solution being prepared at a
7 relatively low temperature; and
8 C. forming a mixture by mixing said joint drink base and said cartilage
9 supplement solution.
- 1 19. A method according to claim 18 comprising the further step of, after step C.
2 D. adjusting the pH of said mixture to be in the range of 2.2 to 4.6.
- 1 20. A method according to claim 16 wherein said cartilage supplement is glucosamine.
- 1 21. A method according to claim 18 wherein said relatively high temperature is above
2 190°F and said relatively low temperature is below 160°F.
- 1 22. A method according to claim 18 wherein said cartilage supplement is selected from
2 glucosamine associated with a counter ion and a glucosamine derivative.
- 1 23. A method according to claim 18 wherein said cartilage supplement is glucosamine
2 hydrochloride.
- 1 24. A method according to claim 18 wherein said steps B and C are performed
2 concurrently.
- 1 25. A method according to claim 19 wherein steps B, C and D are performed
2 concurrently.

- 1 26. A method according to claim 18 comprising the further step of carbonating said
2 mixture.
- 1 27. A method of treating cartilage dysfunction in a human comprising
2 (i) preparing a cartilage enhancing beverage, said beverage containing
3 a mixture of a juice drink base and a cartilage supplement-containing solution
4 wherein:
5 A. said juice drink base is a water-based solution comprising -
6 juice, and prepared using a pasteurization process at a relatively high process
7 temperature, and
8 B. said cartilage-supplement containing solution is a water-based
9 solution comprising one or more cartilage supplements and prepared at a
10 relatively low process temperature.
11 (ii) administering the cartilage enhancing beverage to a human in need
12 of such treatment.
- 1 28. The method according to claim 27 wherein the mixture is formed at a relatively
2 low mixing temperature.
- 1 29. The method according to claim 27 wherein said cartilage supplement is
2 glucosamine hydrochloride.
- 1 30. The method according to claim 28 wherein the relatively high process temperature
2 is above 190°F, the relatively low process temperature is below 160°F, and the mixing
3 temperature is between 40°F and 160°F.
- 1 31. The method of claim 27 wherein the juice drink base selected from the group
2 consisting of water, flavored water, fruit juice concentrate, fruit juice, fruit flavored juice,
3 and combinations thereof.
- 1 32. The method of claim 28 wherein the beverage is carbonated.

1 33. The method of claim 31 wherein the beverage further includes ingredients selected
2 from the group consisting of proteins, fats, carbohydrates, vitamins, minerals and
3 combinations thereof.

1 34. The method of claim 30 wherein the cartilage supplement is glucosamine in
2 association with a counter ion in combination with a cartilage supplement selected from
3 the group consisting of hyaluronic acid, chondroitin, chondroitin sulfate, and
4 combinations thereof.

1 35. A beverage for human consumption comprising:
2 a mixture of a juice drink base and a cartilage enhancing solution wherein:
3 (i) said juice drink base is selected from the group consisting of water,
4 flavored water, fruit juice, fruit flavored juice, and combinations thereof; and
5 (ii) said cartilage enhancing solution comprises glucosamine
6 hydrochloride.

1 36. The beverage according to claim 35 further including ingredients selected from
2 natural sweeteners, artificial sweeteners, preservatives and combinations thereof.

1 37. The beverage according to claim 36 further including ingredients selected from
2 vitamins, minerals, proteins, fats, carbohydrates, and combinations thereof.

- 1 38. A method for preparing a beverage for human consumption comprising:
2 (i) providing a drink base solution;
3 (ii) providing a cartilage enhancing solution comprising one or more
4 cartilage supplements; and
5 (iii) mixing said drink base solution and said cartilage enhancing
6 solution to form the beverage wherein said beverage has between approximately
7 1% to approximately 10% by weight cartilage supplement.
- 1 39. The method according to claim 38 wherein the drink base is selected from water,
2 juice, fruit juice, fruit flavored juice, and combinations thereof.
- 1 40. The method according to claim 38 wherein the cartilage supplement is selected
2 from glucosamine, glucosamine salts, glucosamine derivatives, and combinations thereof.
- 1 41. The method according to claim 40 wherein the cartilage supplement is glucosamine
2 hydrochloride.
- 1 42. The method according to claim 39 wherein the drink base solution further includes
2 sweeteners.
- 1 43. The method according to claim 39 wherein the drink base further includes
2 vitamins.
- 1 44. The method according to claim 39 wherein the drink base further includes
2 minerals.
- 1 45. The method according to claim 38 wherein the mixing step is performed at ambient
2 temperature.
- 1 46. The method according to claim 38 wherein the mixing step is performed at
2 elevated temperature.

1 47. The method according to claim 46 wherein the elevated temperature is above
2 approximately 160°F.

1 48. The method according to claim 38 comprising the further step of, after the mixing
2 step, carbonating the beverage.

1 49. The method according to claim 48 wherein the mixing step is performed at low
2 temperature.

1 50. The method according to claim 49 wherein the low temperature is above
2 approximately 35°F and below ambient temperature.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/40267

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/70, 35/78, 38/00, 59/00; A23L 1/30, 2/00
US CL : 424/600, 195.1; 426/72, 74, 590, 599; 514/2, 23, 25, 54, 62, 492

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/600, 195.1; 426/72, 74, 590, 599; 514/2, 23, 25, 54, 62, 492

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4,766,209 A (CHEN et al.) 23 August 1988 (23.08.1988), column 3, lines 30-38, column 4, lines 65, 68, columns 5, 6.	1, 3, 5, 6, 8-12, 14-16, 38-40, 42, 44, 48-50
X	US 3,679,660 A (MAGNUS) 25 July 1972 (25.07.1972), column 2, lines 29-71.	38, 39, 40, 42, 45
X	STN/CAS, file CAPLUS, Acc. No. 1972:530609, Doc. No. 77:130609, 'Compositions for the treatment of degenerative joint diseases', (DE 2103387 A (LAUSCHER) 17 August 1972 (17.08.1972)), abstract.	1, 3, 5-11, 13-17, 27-31, 35, 36, 38-42, 45
X	US 5,843,919 A (BURGER) 01 December 1998 (01.12.1998), column 3, lines 20-68, column 4, lines 1-63.	35-45

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	"&" document member of the same patent family

Date of the actual completion of the international search

06 October 2000 (06.10.2000)

Date of mailing of the international search report

23 OCT 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Frank Choi

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/40267

Continuation of B. FIELDS SEARCHED Item 3: WEST, STN/CAS

search terms: glucosamine, chondroitin, hyaluronic acid, cartilage, juice, pasteurization, drink, beverage